

Who receives a medical evaluation for infertility in the United States?

Leslie V. Farland, Sc.M.,^a Ai-ris Y. Collier, M.D.,^b Katharine F. Correia, M.P.H.,^c Francine Grodstein, Sc.D.,^{a,d} Jorge E. Chavarro, M.D., Sc.D.,^{a,d,e} Janet Rich-Edwards, Sc.D.,^{a,d,f} and Stacey A. Missmer, Sc.D.^{a,d,g}

^a Department of Epidemiology, Harvard T.H. Chan School of Public Health; ^b Department of Obstetrics and Gynecology, Beth Israel Deaconess Medical Center; ^c Department of Biostatistics, Harvard T.H. Chan School of Public Health;

^d Channing Division of Network Medicine, Brigham and Women's Hospital and Harvard Medical School; ^e Department of Nutrition, Harvard T.H. Chan School of Public Health; ^f Connors Center for Women's Health and Gender Biology, Brigham and Women's Hospital and Harvard Medical School; and ^g Department of Obstetrics, Gynecology, and Reproductive Biology, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts

Objective: To investigate characteristics of receiving a medical evaluation for infertility among infertile women.

Design: Prospective cohort.

Setting: Academic institution.

Patient(s): A total of 7,422 women who reported incident infertility between 1989 and 2009 in the Nurses' Health Study II.

Intervention(s): None.

Main Outcome Measure(s): Report of receiving a medical evaluation for infertility.

Result(s): Approximately 65% of women who reported infertility had a medical evaluation for infertility. Infertile women who were parous (relative risk [RR] = 0.81, 95% confidence interval [CI] 0.78–0.84), older, current smokers (RR = 0.89, 95% CI 0.83–0.96), or who had a higher body mass index (BMI) were less likely to report receiving a medical infertility evaluation. Infertile women who exercised frequently, took multivitamins (RR = 1.03, 95% CI 1.00–1.07), lived in states with comprehensive insurance coverage (RR = 1.09, 95% CI 1.00–1.19), had a high household income, or who had a recent physical examination (RR = 1.15, 95% CI 1.06–1.24) were more likely to report receiving a medical infertility evaluation.

Conclusion(s): These findings highlight demographic, lifestyle, and access barriers to receiving medical infertility care. Historically, the discussion of barriers to infertility care has centered on financial access, geographic access, and socioeconomic status. Our findings build off literature by supporting previously reported associations and showcasing the importance of demographic and lifestyle factors in accessing care. (Fertil Steril® 2016;105:1274–80. ©2016 by American Society for Reproductive Medicine.)

Key Words: Infertility, fertility evaluation, fertility treatment, barriers to access, barriers to care

Discuss: You can discuss this article with its authors and with other ASRM members at <http://fertsterfforum.com/farland1-medical-evaluation-infertility-us/>



Use your smartphone to scan this QR code and connect to the discussion forum for this article now.*

* Download a free QR code scanner by searching for "QR scanner" in your smartphone's app store or app marketplace.

In the United States from 2006–2010, approximately 1.5 million couples report being affected by infertility each year (1, 2). Understanding the burden of this disease has become a national priority (3), with the U.S. Department of Health and Human Services and Centers for Disease Control and Prevention releasing a National

Public Health Action Plan for the Detection, Prevention, and Management of Infertility (4). The National Survey for Family Growth (NSFG), estimated that among women with fertility problems, in 1982, 1995, and 2006–2010, only 41%–46% of women have ever used any type of infertility service (including both medical help to get pregnant and

to prevent miscarriage) and this proportion has not varied across time (5).

Data from national surveys and clinic-based studies investigating barriers in access to infertility care have been limited in their scope, focusing on differences by race, age, cause of infertility, and socioeconomic factors (1, 5–15). The most consistently investigated predictors for accessing fertility care are financial access (insurance coverage, income, and high educational attainment) and white race. In addition to these factors, healthy behaviors, lifestyle factors, and access to the medical system may contribute to whether or not an infertile couple has a medical evaluation for their condition. However, these factors have not been thoroughly investigated. The

Received October 5, 2015; revised December 18, 2015; accepted December 22, 2015; published online January 16, 2016.

L.V.F. has nothing to disclose. A.Y.C. has nothing to disclose. K.F.C. has nothing to disclose. F.G. has nothing to disclose. J.E.C. has nothing to disclose. J.R.-E. has nothing to disclose. S.A.M. has nothing to disclose.

Supported by the National Institutes of Health (HD59955 and UM1 CA176726; and T32HD060454 [to L.V.F.]).

Reprint requests: Leslie V. Farland, Sc.M., Harvard T.H. Chan School of Public Health, Department of Epidemiology, 677 Huntington Avenue, Boston, Massachusetts 02115 (E-mail: lfarland@hspf.harvard.edu).

current research may have methodological limitations due to small sample sizes, poor response rates, and using cross-sectional study designs that could lead to reverse causation or recall bias.

Using data from the Nurses' Health Study II, a large prospective cohort of female medical professionals, we evaluated a broad range of characteristics associated with utilization of fertility evaluation. We hypothesized that previously investigated characteristics including age, race/ethnicity, income, and insurance coverage would alter fertility care utilization. We also hypothesized that previously uninvestigated lifestyle characteristics such as body mass index (BMI), cigarette smoking, vitamin use, exercise, and routine physical examination history would influence whether or not women have a medical evaluation for infertility.

MATERIALS AND METHODS

The Nurses' Health Study II is a prospective cohort study that began in 1989 when 116,430 registered nurses, 25–42 years old, returned a mailed questionnaire regarding their health and lifestyle. At recruitment, women lived in 1 of 14 states. However, the participants have since moved to all 50 states. Follow-up questionnaires are sent biennially, with a cumulative follow-up rate from the original cohort of $\geq 90\%$. Informed consent was obtained from all participants and the study was approved by the Institutional Review Board of Brigham and Women's Hospital. For the current analysis, premenopausal women with no history of hysterectomy, oophorectomy, or tubal ligation were followed from 1989, when the cohort began, through 2009.

Collection of Information on Fertility Evaluation

To define infertility status, women were asked to self-report if they had "tried to become pregnant for more than 1 year without success" on every questionnaire cycle from 1989–2001, and in 2005 and 2009. We restricted our study population to women who reported incident infertility after the first questionnaire cycle ($n = 7,422$). To define our outcome, women were then asked what was the cause for their infertility and were given the following choices: "not investigated, not found, tubal blockage, ovulatory disorder, endometriosis, cervical mucus factors, spousal infertility, and/or other." Women could report multiple causes for infertility. Women who reported "not investigated" were considered not to have sought infertility evaluation ($n = 2,598$). Women who did not report "not investigated" but instead reported a cause for infertility or that the cause was "not found" were classified as having reported medical evaluation and diagnosis of infertility ($n = 4,824$).

Reliability and Validity of Self-Reported Infertility

Although validation data were not available on all types of infertility, a validation study of self-reported ovulatory disorder infertility was conducted among a random subset of 100 women in the Nurses' Health Study II who cited ovulatory infertility as a physician identified infertility cause on the questionnaire. More than 93% of the women who responded

to the supplemental questionnaire reported diagnostic test results and/or indicative treatment for ovulatory infertility indicating that a conventional infertility workup was performed. In addition, among a subsample of the original 100 women, 40 random women whose participant medical records were reviewed, 95% of these women had indication in their medical records (diagnostic test and/or treatment) confirming medically diagnosed ovulatory disorder infertility and a conventional infertility workup (16). We also see high validity of our measure of self-reported fertility treatment in this cohort across time ($\geq 84\%$ concordance) and with medical records (74% of medical records confirmed women's reported treatment, whereas the remaining records generally contained no information on specific treatments) (17). These validation estimates verify the nurses' ability to accurately report their experience with conventional physician-based infertility evaluation.

Covariates

Because we wished to prospectively quantify the association between the self-reported woman's characteristics and her likelihood of seeking fertility evaluation to reduce reverse causation and recall bias, the covariate values were defined approximately 2 years before the first report of infertility, with the exception of marriage, which was collected at time of reported infertility. Demographic factors including age (categorized according to the Society of Assisted Reproductive Technology [SART] age guidelines), race, marriage, and male partner's education in 1999 were analyzed. We considered reproductive characteristics including nulliparity, history of uterine fibroids, history of endometriosis, and history of spontaneous abortion. We also considered several self-reported lifestyle factors, including current BMI (in kilograms per meter squared) based on weight and height measurements as discussed in detail previously (18), BMI at age 18 years, cigarette smoking status, alcohol intake, current multivitamin use, and physical activity (measured in metabolic equivalent of task hours/week based on weekly reported recreational physical activity) (19). Finally, we considered factors related to access, including at least one routine physical examination, physical examination for general health symptoms, and annual household income. State-mandated insurance coverage of fertility treatment was defined as "comprehensive coverage" (state-mandated infertility treatment coverage including assisted reproductive technology [ART]), "limited coverage" (state-mandated infertility coverage that included diagnosis and treatment but may exclude IVF treatment or did not specify treatment coverage), and "offer only" (state-mandated offer of an insurance policy that includes fertility treatments available for purchase) based on history of state legislature on fertility treatment, which was updated at each questionnaire cycle (11, 17, 20). All models were mutually adjusted for other demographic, lifestyle, and access covariates.

Data Analysis

Log-binomial models were used to estimate relative risks and confidence intervals of seeking an infertility evaluation (21).

In a few instances, the models did not converge and log-Poisson models, which provide consistent but not fully efficient estimates of the relative risk and its confidence intervals, were used (22). To test for linear trend, variables were set to mean values within each category and treated as linear. Sensitivity analyses expanded the outcome definition to include women who reported infertility evaluations on a later questionnaire cycle than the reported experience of infertility.

RESULTS

Of the 7,422 women reporting incident infertility, approximately 65% reported a medical infertility evaluation (Table 1). The mean age at first report of infertility was 35.1 years ($SD = 4.7$) among those who reported medical evaluation for infertility and 36.2 years ($SD = 4.7$) among those who did not. Of the women who reported medical evaluation for infertility, 9% reported tubal infertility as a possible infertility cause, 27% reported ovulatory dysfunction, 12% reported endometriosis, 4% reported cervical mucosal factors, 19% reported male factor infertility, 19% reported “other” reason, and 33% reported that the reason for infertility was not found.

Several demographic characteristics were related to fertility evaluation among women with infertility. Infertile women of older age were less likely to report a fertility evaluation compared with their younger peers ($P < .001$, test for linear trend) (Table 2). Infertile women whose male partners had graduate level education were more likely to report an evaluation than women whose partners had attained less than a 4-year college degree ($P < .001$, test for linear trend). No significant difference was observed by race or marital status. Among reproductive factors, infertile women who were parous were less likely to report a medical evaluation compared with nulliparous infertile women (relative risk [RR] = 0.81, 95% confidence interval [CI] 0.78–0.84). Whereas infertile women who ever reported ultrasound or hysterectomy-diagnosed uterine fibroids were not more or less likely to report a medical evaluation. Those with a history of surgically confirmed endometriosis were more likely to report having a medical evaluation than infertile women without a prior endometriosis diagnosis (RR = 1.27, 95% CI 1.20–1.35). History of spontaneous abortion was not associated with evaluation.

Among lifestyle factors, although there was no association between BMI at age 18 years and fertility evaluation ($P = .16$, test for linear trend), there were significant findings among other lifestyle covariates (Table 3). There was a significant inverse relationship between adult BMI and likelihood of reporting a fertility evaluation among infertile women ($P = .01$, test for linear trend). Infertile women who were current smokers were less likely to report having a medical evaluation than never smokers (RR = 0.89, 95% CI 0.83–0.96), whereas alcohol intake was unrelated to infertility diagnosis. In addition, infertile women who exercised frequently were more likely report a fertility evaluation than their sedentary counterparts ($P = .04$, test for linear trend). Compared with infertile women who did not use multivitamins, those who

TABLE 1

Age-standardized characteristics of women by whether they sought a medical evaluation for infertility: Nurses' Health Study II (1989–2009).

Characteristic	Medical evaluation for infertility	
	No (n = 2,598)	Yes (n = 4,824)
Age at infertility (y) ^a	36.19 (4.72)	35.14 (4.70)
Race		
White	91	91
Nulliparous		
Yes	47	61
State of residence with mandated insurance coverage for infertility treatment		
No coverage	56	50
Comprehensive coverage	2	3
Limited coverage	18	20
Offer only	24	27
BMI (kg/m ²)		
<18.5	4	4
18.5–24.9	62	65
25–29.9	20	18
30 ⁺	15	12
Annual household income (\$)		
<50,000	16	13
50,000–99,999	49	49
100,000–149,999	23	22
>150,000	12	16
Partner's highest level of education		
<4-y college	35	30
4-y college	30	32
Graduate school	30	33
Recent physical examination		
No	11	8
Yes, for symptoms	23	25
Yes, for screening	66	68
Smoking status		
Never smoker	68	69
Past smoker	20	22
Current smoker	12	9
Exercise, MET (h/wk)		
<3	15	13
3–8.9	22	21
9–17.9	22	22
>18	41	45
Multivitamin use		
Yes	53	57
No	47	43
Year of first infertility report		
1989–1992	51	51
1993–1996	32	30
1997–2009	18	19
Fertility diagnosis among those who had a medical evaluation ^b		
Tubal		9
Ovulatory dysfunction		27
Endometriosis		12
Cervical mucosal factors		4
Male factor infertility		19
Other reason		19
Reason not found		33

Note: Values are means (SD) or percentages and are standardized to the age distribution of the study population. Values of polytomous variables may not sum to 100% due to rounding. BMI = body mass index; MET = metabolic equivalent of task.

^a Value is presented as mean (SD) and not age adjusted.

^b Categories not mutually exclusive.

Farland. Predictors of infertility evaluation. *Fertil Steril* 2016.

TABLE 2

Demographic factors and the risk of seeking an infertility evaluation in the Nurses' Health Study II, 1989–2009.

Demographic factor	N (% infertility evaluation)	Crude	Adjusted
		Risk ratio (95% confidence interval)	
Age (y)			
<35	2,306 (69.3)	1.00 (Referent)	1.00 (Referent)
35–37	991 (62.8)	0.91 (0.87–0.95)	0.92 (0.88–0.97)
38–40	621 (64.8)	0.93 (0.89–0.98)	0.93 (0.88–0.98)
≥40	906 (58.1)	0.84 (0.80–0.88)	0.85 (0.81–0.89)
P value, test for trend ^a		<.001	<.001
Race			
White	4,412 (65.2)	1.00 (Referent)	1.00 (Referent)
Black	68 (58.1)	0.89 (0.76–1.04)	0.91 (0.79–1.06)
Asian	141 (68.4)	1.05 (0.95–1.15)	1.02 (0.93–1.12)
Other	203 (60.2)	0.92 (0.85–1.01)	0.93 (0.85–1.02)
Marital status			
Never married	539 (67.5)	1.00 (Referent)	1.00 (Referent)
Married/domestic partnership	3,458 (64.7)	0.96 (0.91–1.01)	1.04 (0.97–1.11)
Divorced/separated/widowed	246 (59.1)	0.88 (0.80–0.96)	0.99 (0.91–1.08)
Partner's education			
<4-y college	1,286 (61.2)	1.00 (Referent)	1.00 (Referent)
4-y college	1,394 (66.3)	1.08 (1.04–1.14)	1.05 (1.01–1.10)
Graduate school	1,436 (66.6)	1.09 (1.04–1.14)	1.07 (1.02–1.12)
P value, test for trend		<.001	<.001
Reproductive factors			
Nulliparous			
Yes	2,731 (71.1)	1.00 (Referent)	1.00 (Referent)
No	1,711 (56.7)	0.80 (0.77–0.83)	0.81 (0.78–0.84)
Ever uterine fibroids			
No	4,527 (64.9)	1.00 (Referent)	1.00 (Referent)
Yes	297 (66.4)	1.02 (0.96–1.10)	1.06 (0.99–1.14)
Ever endometriosis			
No	4,595 (64.3)	1.00 (Referent)	1.00 (Referent)
Yes	229 (84.2)	1.31 (1.24–1.38)	1.27 (1.20–1.35)
Ever spontaneous abortion			
No	4,101 (65.5)	1.00 (Referent)	1.00 (Referent)
Yes	522 (61.7)	1.01 (0.93–1.10)	0.93 (0.85–1.01)

Note: Log binomial regression adjusted for age, race, nulliparity, insurance coverage in state of residence, sexual identification, marital status, living arrangement, income, husband's education level, recent physical examination, body mass index (BMI), smoking status, exercise, multivitamin use, spontaneous abortion, and recent illness or surgery.

^a Wald test used to assess linear trend among categories.

Farland. Predictors of infertility evaluation. *Fertil Steril* 2016.

did were more likely to report an evaluation (RR = 1.03, 95% CI 1.00–1.07).

Last, infertile women who resided in states with any level of state-mandated insurance coverage for infertility treatment were more likely to report having an infertility evaluation compared with those residing in states without mandated insurance coverage (offer only, RR = 1.07, 95% CI 1.03–1.11; limited coverage, RR = 1.09, 95% CI 1.04–1.14; comprehensive coverage, RR = 1.09, 95% CI 1.00–1.19) (Table 4). Infertile women with higher household incomes were more likely to report having an infertility evaluation ($P=.05$, test for linear trend). Compared with infertile women who had not reported a physical examination before their infertility, women who had a recent (~2 years before infertility) general physical examination (RR = 1.14, 95% CI 1.06–1.22) or an examination for symptoms of any health condition (RR = 1.15, 95% CI 1.06–1.24) were more likely to report having a fertility evaluation. In sensitivity analyses, which expanded the outcome definition to include delayed report of fertility investigation, overall trends did not significantly change.

DISCUSSION

Infertile women who reported receiving a medical infertility evaluation were different from their peers with uninvestigated infertility on several important demographic, lifestyle, and access characteristics. Among this cohort of registered nurses who reported infertility, most women (65%) reported receiving a medical diagnosis of their fertility. To our knowledge this is the largest prospective study with inclusion of detailed confounding control that investigates a range of predictors of medical evaluation for infertility.

The NSFG has collected nationally representative estimates for infertility service use (both medical help getting pregnant and help preventing miscarriage) for women aged 15–44 years since 1982 (5, 9, 15). Although they report no significant difference across time in infertility service utilization for help getting pregnant among women with infertility problems, nulliparous women with infertility problems were less likely to use services in 2006–2010 (38%) compared with 1982 (56%) (5). This may reflect societal shifts in delays in age at conception of first child

TABLE 3

Lifestyle factors and the risk of seeking an infertility evaluation in the Nurses' Health Study II, 1989–2009.

Lifestyle factor	N (% infertility evaluation)	Crude	Adjusted
		Risk ratio (95% confidence interval)	
BMI at age 18 (kg/m ²) ^a			
<18.5	827 (66.3)	1.02 (0.98–1.07)	1.02 (0.98–1.07)
18.5–24.9	3,411 (64.8)	1.00 (Referent)	1.00 (Referent)
25.5–29.9	378 (64.8)	1.00 (0.94–1.07)	1.00 (0.94–1.07)
+30	151 (61.4)	0.95 (0.86–1.05)	0.95 (0.86–1.05)
P value, test for trend ^b		.13	.16
BMI (kg/m ²)			
<18.5	196 (69.5)	1.05 (0.97–1.14)	1.05 (0.97–1.13)
18.5–24.9	2,880 (65.9)	1.00 (Referent)	1.00 (Referent)
25.5–29.9	808 (62.9)	0.95 (0.91–1.00)	0.99 (0.95–1.04)
30+	529 (59.2)	0.90 (0.85–0.95)	0.96 (0.90–1.01)
P value, test for trend		.02	.01
Smoking status			
Never smoker	3,338 (65.6)	1.00 (Referent)	1.00 (Referent)
Past smoker	1,026 (65.9)	1.00 (0.96–1.05)	1.04 (1.00–1.09)
Current smoker	437 (58.8)	0.90 (0.84–0.95)	0.89 (0.83–0.96)
Alcohol intake, g/d			
No alcohol	1,385 (63.6)	1.00 (Referent)	1.00 (Referent)
<5	1,772 (65.7)	1.06 (1.00–1.13)	0.98 (0.94–1.02)
5.01–10	522 (67.5)	1.03 (0.99–1.08)	1.00 (0.94–1.06)
>10	421 (62.0)	0.97 (0.91–1.04)	0.94 (0.88–1.00)
P value, test for trend		.69	.12
Exercise, MET (h/wk)			
<3	542 (60.4)	1.00 (Referent)	1.00 (Referent)
3–8.9	895 (63.2)	1.05 (0.98–1.12)	1.02 (0.96–1.09)
9–17.9	951 (65.0)	1.08 (1.01–1.15)	1.04 (0.97–1.10)
18–26.9	581 (65.6)	1.09 (1.01–1.17)	1.05 (0.98–1.13)
27–41.9	628 (66.5)	1.10 (1.03–1.18)	1.05 (0.98–1.12)
+42	751 (68.3)	1.13 (1.06–1.21)	1.07 (1.00–1.14)
P value, test for trend		<.001	.04
Multivitamin use			
No	1,858 (63.2)	1.00 (Referent)	1.00 (Referent)
Yes	2,383 (65.9)	1.04 (1.01–1.08)	1.03 (1.00–1.07)

Note: Log binomial regression adjusted for age, race, nulliparity, insurance coverage in state of residence, sexual identification, marital status, living arrangement, income, husband's education level, recent physical examination, body mass index, smoking status, exercise, multivitamin use, spontaneous abortion, and recent illness or surgery. BMI = body mass index; MET = metabolic equivalent of task.

^a Adjusted for race/ethnicity.

^b Wald test used to assess linear trend among categories.

Farland. Predictors of infertility evaluation. *Fertil Steril* 2016.

and use of services beyond the age of 44 years, which are not captured in the NSFG. In our study population, women were eligible to enter the analysis until menopause, and we observed no difference in utilization patterns after adjusting for age (Table 1).

Our study was consistent with previous work during this time period and found that after adjusting for confounding, "demographic characteristics" (age at infertility, parity, husband's education) and "access characteristics" (income, connection with the medical system, and insurance status) were significant predictors of reporting having received an infertility evaluation (1,5–13, 15,23–25). Being older was predictive of not reporting having had an evaluation to a similar extent as having been previously parous. In previous

studies from this cohort, women who were older were also less likely to use advanced levels of fertility treatment (17). In addition, partners' education level was predictive of being more likely to report having received an infertility evaluation, which is consistent with the previous literature that found couples with high education are more likely to seek care as it may stand in as a marker for socioeconomic status (5, 7–9, 11, 12, 15). Access also played an important role in whether individuals reported receiving a medical diagnosis. Not surprisingly, having financial means, insurance coverage, and connection with the medical system (recent physical examination) were all predictive of reporting a medical evaluation for infertility in our cohort. We did not find significant differences between race and marital status and fertility evaluation, which have been reported previously (5, 7, 11, 14, 26). This may be a reflection of our highly educated, relatively homogenous population among which there may be limited power to detect such differences (black women, n = 68) or this may support previous work in which, after adjustment, the effect of race was attenuated by other socioeconomic factors (9, 15). Historically, the discussion of barriers to infertility care has centered on financial access, geographic access, and socioeconomic status (5–7, 9, 12, 15, 27). However, our findings build off of previous literature by showcasing how lifestyle, demographic, and access factors independent of one another are associated with reporting having received an evaluation for infertility.

Our analyses found that independent of markers of access, there are several predictors of reporting receipt of an infertility evaluation related to lifestyle characteristics. When evaluating the association between lifestyle factors and fertility evaluation, a consistent pattern emerged. Infertile women who reported behaviors that are generally regarded as "healthy lifestyle" were more likely to report having had an evaluation than their peers who exhibited less "healthy lifestyle" behaviors. For example, women who had a lower BMI preceding infertility, who never smoked, who used multivitamins, and who exercised regularly were more likely to have reported having received an infertility diagnosis compared with their less healthy counterparts independent of all other characteristics.

When looking at reproductive health factors that could represent an individual's past engagement with the reproductive medical system (uterine fibroids diagnosis, endometriosis diagnosis, and spontaneous abortion) no clear pattern emerged. Those who had ever been diagnosed with endometriosis were more likely to reported having received a medical evaluation for their infertility 2 years later. However, those with a uterine fibroid diagnosis and those with a spontaneous abortion were not statistically more likely to report having received an evaluation. The former may be confounded by indication, given the strong correlation between infertility and endometriosis.

The findings of this work, specifically the associations with demographic and healthy lifestyle factors, may help guide clinicians who serve as first-line providers for many women who are suffering from uninvestigated infertility. There are important barriers to using infertility care that are

TABLE 4

Access factors and the risk of seeking an infertility evaluation in the Nurses' Health Study II, 1989–2009.

Access factor	N (% infertility evaluation)	Crude	Adjusted
		Risk ratio (95% confidence interval)	
Mandated insurance coverage			
No coverage	2,452 (63.6)	1.00 (Referent)	1.00 (Referent)
Offer only	1,274 (66.7)	1.05 (1.01–1.09)	1.07 (1.03–1.11)
Limited coverage	953 (66.0)	1.04 (0.99–1.08)	1.09 (1.04–1.14)
Comprehensive coverage	145 (69.0)	1.09 (0.99–1.19)	1.09 (1.00–1.19)
Household annual income, \$			
<50,000	377 (59.1)	0.94 (0.87–1.01)	0.94 (0.87–1.00)
50,000–99	1,414 (63.0)	1.00 (Referent)	1.00 (Referent)
100,000–149	647 (62.5)	0.99 (0.94–1.05)	0.97 (0.92–1.03)
>150,000	469 (68.7)	1.09 (1.03–1.16)	1.03 (0.97–1.09)
<i>P</i> value, test for trend ^b		<.001	.05
Recent physical examination ^a			
No	349 (56.5)	1.00 (Referent)	1.00 (Referent)
Yes for symptoms	1,110 (65.8)	1.17 (1.09–1.25)	1.15 (1.06–1.24)
Yes for screening	3,081 (65.9)	1.17 (1.08–1.26)	1.14 (1.06–1.22)
Recent major illness or surgery			
No	4,085 (64.7)	1.00 (Referent)	1.00 (Referent)
Yes	739 (66.8)	1.03 (0.99–1.08)	1.01 (0.97–1.06)

Note: Log binomial regression adjusted for age, race, nulliparity, insurance coverage in state of residence, sexual identification, marital status, living arrangement, income, husband's education level, recent physical examination, body mass index (BMI), smoking status, exercise, multivitamin use, spontaneous abortion, and recent illness or surgery.

^a This includes physical examinations outside of an infertility work-up, measured 2 years before reported infertility.

^b Wald test used to assess linear trend among categories.

Farland. Predictors of infertility evaluation. *Fertil Steril* 2016.

not fully addressed by differences in financial access and insurance coverage. In our data we are unable to determine whether these differences were a product of referral patterns or differences in seeking access by the individual. Therefore, these barriers need to be thoughtfully considered by doctors administering care to reproductive aged women and may be targeted for public health interventions encouraging women to seek medical evaluation for their infertility.

This study also highlights potential methodological limitations of research restricted to women enrolled from a fertility clinic setting. Our findings suggest that associations seen within studies using only infertility clinic populations should be interpreted carefully. If couples who present in the fertility clinic setting exhibit healthier lifestyle characteristics (lower BMI, multivitamin use, regular exercise) in addition to markers of higher socioeconomic status (higher household income, higher education) than their infertile peers who do not receive a medical evaluation, then results from studies restricted to fertility clinic practices may not be generalizable to the broader population of women experiencing infertility. In addition, depending on the causal question of interest, using a population restricted to fertility clinic patients may result in selection bias.

As with all studies, there are limitations of this current research. Covariates and outcomes of interest were based on self-report, which may cause concern about the potential for misclassification. In addition, given the health knowledge of our cohort, there is potential for misclassification of report of fertility evaluation. Given validation evidence and the specificity of our questions, we are making the assumption that all participants who reported a specific infertility cause and who also did not report that their infertility was "not

investigated," in fact had their fertility investigated in a medical setting. Although we were not able to validate reports on all types of infertility, our validation studies on report of ovulatory infertility and fertility treatment provided evidence that misclassification of the participant's infertility experience is unlikely. Indeed, among women reporting ovulatory infertility there is high concordance (95%) with medical records, indicating that nearly all women who reported ovulatory infertility had a diagnosis and medical evaluation for infertility. In addition, if misclassification of our outcome did exist, we would expect any possible misclassification to be nondifferential (i.e., random with respect to our exposures) and thus would attenuate our results toward the null, which would thus lead to underestimation of true effects.

The population under study was a group of medical professionals and was relatively homogenous in terms of race and education level. In fact, the distribution of infertility differs slightly from previous reports among the population that utilizes ART in the United States (28) and the proportion of women who reported having received a fertility evaluation was higher (65%) than estimates previously reported from the NSFG, which estimated that among women with fertility problems, in 2006–2010, 36% of women reported having ever used medical help to get pregnant, this prevalence has not statistically significantly changed over time among NSFG cross-sectional sample populations (5). These results represent a group with high medical knowledge and access, among whom we still observed meaningful disparities in reporting receiving an infertility evaluation. We hypothesize that these differences in access would be even greater among the less homogeneous general population of women experiencing infertility.

Despite these limitations, this study has several strengths including its large sample size, detailed temporal evaluation of predictors and confounders, high response rate, and validated exposure and outcome measures. Our analysis used log binomial regression to quantify relative risks, as opposed to logistic regression to quantify odds ratios, which are known to overestimate associations when the outcome is common (29).

In conclusion, the Centers of Disease Control and Prevention has recently stated that the detection of infertility and management of infertility care is a national public health priority (3, 4). This research furthers our understanding of the wide range of barriers to infertility care and may be used to guide proposed scientific and programmatic projects to encourage increased medical evaluation for infertility. The Public Health Action Plan calls for more research in the detection of infertility, prevention of infertility, and management of infertility including better understanding the safety and efficacy of infertility treatments (4). The findings from our current work highlight important issues of external and internal validity that may arise if future research is conducted solely within infertility clinic settings. Understanding the many facets of infertility is a newly stated national priority in the United States and the findings of this study help to elucidate our understanding of the utilization of infertility care.

REFERENCES

- Chandra A, Copen CE, Stephen EH. Infertility and impaired fecundity in the United States, 1982–2010: data from the National Survey of Family Growth. In: National health statistics reports. 67th ed. Hyattsville, MD: National Center for Health Statistics. Available at: <http://www.cdc.gov/nchs/data/nhsr/nhsr067.pdf>; 2013. Accessed January 19, 2016.
- Thoma ME, McLain AC, Louis JF, King RB, Trumble AC, Sundaram R, et al. Prevalence of infertility in the United States as estimated by the current duration approach and a traditional constructed approach. *Fertil Steril* 2013;99: 1324–31.e1.
- Macaluso M, Wright-Schnapp TJ, Chandra A, Johnson R, Satterwhite CL, Pulver A, et al. A public health focus on infertility prevention, detection, and management. *Fertil Steril* 2010;93:16.e1–10.
- National Public Health Action Plan for detection, prevention, and management of infertility. Atlanta, GA: Centers of Disease Control and Prevention; 2014.
- Chandra A, Copen CE, Stephen EH. Infertility service use in the United States: data from the national survey of family growth, 1982–2010. *Natl Health Stat Rep* 2014;73:1–21.
- Farley Ordovensky Staniec J, Webb NJ. Utilization of infertility services: how much does money matter? *Health Serv Res* 2007;42:971–89.
- Jain T. Socioeconomic and racial disparities among infertility patients seeking care. *Fertil Steril* 2006;85:876–81.
- Wilcox LS, Mosher WD. Use of infertility services in the United States. *Obstet Gynecol* 1993;82:122–7.
- Stephen EH, Chandra A. Use of infertility services in the United States: 1995. *Fam Plann Perspect* 2000;32:132–7.
- Hirsch MB, Mosher WD. Characteristics of infertile women in the United States and their use of infertility services. *Fertil Steril* 1987;47:618–25.
- Bitler M, Schmidt L. Health disparities and infertility: impacts of state-level insurance mandates. *Fertil Steril* 2006;85:858–65.
- Jain T, Hornstein MD. Disparities in access to infertility services in a state with mandated insurance coverage. *Fertil Steril* 2005;84:221–3.
- Peterson MM. Assisted reproductive technologies and equity of access issues. *J Med Ethics* 2005;31:280–5.
- Greil AL, McQuillan J, Shreffler KM, Johnson KM, Slauson-Blevins KS. Race-ethnicity and medical services for infertility: stratified reproduction in a population-based sample of U.S. women. *J Health Soc Behav* 2011;52: 493–509.
- Chandra A, Stephen EH. Infertility service use among U.S. women: 1995 and 2002. *Fertil Steril* 2010;93:725–36.
- Rich-Edwards JW, Goldman MB, Willett WC, Hunter DJ, Stampfer MJ, Colditz GA, et al. Adolescent body mass index and infertility caused by ovulatory disorder. *Am J Obstet Gynecol* 1994;171:171–7.
- Farland LV, Missmer SA, Rich-Edwards J, Chavarro JE, Barbieri RL, Grodstein F. Use of fertility treatment modalities in a large United States cohort of professional women. *Fertil Steril* 2014;101:1705–10.
- Manson JE, Willett WC, Stampfer MJ, Colditz GA, Hunter DJ, Hankinson SE, et al. Body weight and mortality among women. *N Engl J Med* 1995;333: 677–85.
- Wolf AM, Hunter DJ, Colditz GA, Manson JE, Stampfer MJ, Corsano KA, et al. Reproducibility and validity of a self-administered physical activity questionnaire. *Int J Epidemiol* 1994;23:991–9.
- Henne MB, Bendorf MK. Insurance mandates and trends in infertility treatments. *Fertil Steril* 2008;89:66–73.
- Wacholder S. Binomial regression in GLIM: estimating risk ratios and risk differences. *Am J Epidemiol* 1986;123:174–84.
- Zou GA. modified Poisson regression approach to prospective studies with binary data. *Am J Epidemiol* 2004;159:702–6.
- Abusief ME, Missmer SA, Barbieri RL, Jain T, Hornstein MD. Geographic distribution of reproductive endocrinology and infertility (REI) fellowships in the United States. *Fertil Steril* 2009;91:1636–41.
- Smith JF, Eisenberg ML, Glidden D, Millstein SG, Cedars M, Walsh TJ, et al. Socioeconomic disparities in the use and success of fertility treatments: analysis of data from a prospective cohort in the United States. *Fertil Steril* 2011; 96:95–101.
- Mosher WD, Bachrach CA. Understanding U.S. fertility: continuity and change in the National Survey of Family Growth, 1988–1995. *Fam Plann Perspect* 1996;28:4–12.
- White L, McQuillan J, Greil AL. Explaining disparities in treatment seeking: the case of infertility. *Fertil Steril* 2006;85:853–7.
- Feinberg EC, Larsen FW, Catherino WH, Zhang J, Armstrong AY. Comparison of assisted reproductive technology utilization and outcomes between Caucasian and African American patients in an equal-access-to-care setting. *Fertil Steril* 2006;85:888–94.
- Centers for Disease Control and Prevention. ASRM, Society for Assisted Reproductive Technology. 2011 Assisted Reproductive Technology Fertility Clinic Success Rates Report. Atlanta, GA: US Dept of Health and Human Services; 2011.
- Davies HT, Crombie IK, Tavakoli M. When can odds ratios mislead? *BMJ* 1998;316:989–91.