



## COMMENTARY

# SARS-CoV-2 and the testis: similarity with other viruses and routes of infection



Walter D. Cardona Maya<sup>1</sup>, Stefan S. Du Plessis<sup>2,3</sup>, Paula A. Velilla<sup>4</sup>

## ABSTRACT

Since the start of the latest coronavirus (SARS-CoV-2) outbreak, the number of infected individuals and cases of coronavirus disease (COVID-19) has been increasing exponentially worldwide. Of interest is existing evidence that orchitis can develop due to coronavirus infection. It is therefore not unreasonable to believe that SARS-CoV-2 could be transmitted by semen. Consequently, it is of paramount importance that individuals who could potentially be infected take all possible care to mitigate the likely risk of passing on the infection through sexual intercourse.

**S**ince the novel coronavirus (SARS-CoV-2) outbreak in December 2019, the number of cases of infected individuals has been increasing exponentially, leading to the coronavirus disease (COVID-19) being declared a global pandemic on the 11 March 2020.

Rothe and collaborators (Rothe *et al.* 2020) were the first to report that a male acquired the infection through transmission that appears to have occurred during the incubation period of the index patient, who did not present with any symptoms at that stage. This case raises questions about possible new routes of transmission.

There is evidence about the relationship between members of the coronavirus family and orchitis in both humans (SARS-CoV) (Xu *et al.* 2006) and cats (feline coronavirus) (Sigurðardóttir *et al.* 2001). However, whether the virus is present in the testicular epithelium remains a matter of debate due to

conflicting results (Zhao *et al.* 2003; Ding *et al.* 2004; Gu *et al.* 2005; Xu *et al.* 2006). As testicular damage and germ cell destruction was clearly observed in these cases (Xu *et al.* 2006), it is not unreasonable to believe that SARS-CoV-2 could be transmitted by semen, potentially providing a possible route of transmission of the pathogen. This could occur in asymptomatic patients, in patients with mild symptoms or even during the incubation period. This potential route of transmission has been proposed for other viral infections that were previously considered unlikely to be transferred sexually, including Ebola (Cardona-Maya *et al.* 2019) and Zika viruses (Cardona Maya *et al.* 2016, Feldmann 2018). It was also recently reported that angiotensin-converting enzyme 2 (ACE2) has been shown to be one of the major receptors that mediate the entry of SARS-CoV-2 into human cells. Using online datasets, Fan and colleagues (Fan *et al.* 2020) clearly demonstrated that ACE2 is highly expressed in Leydig cells and cells of

the seminiferous tubules in the human testis. Thus, the binding of the virus to these ACE2-positive cells could not only cause severe alteration of testicular tissue but also provide a sites for viral infection. It should therefore be of some concern that this recent coronavirus outbreak may have a serious impact on fertility worldwide. Although the actual situation has yet to be substantiated in patients infected with SARS-CoV-2, the American Society for Reproductive Medicine (ASRM) and the Society for Assisted reproductive Technology (SART) have already issued warnings to this extent (SART, 2020). The advice given is that prospective parents, ART patients, gamete donors and gestational carriers who meet the diagnostic criteria for SARS-CoV-2 should avoid becoming pregnant or participate in any fertility programs (Society for Assisted Reproductive Technology, 2020). The Center for Diseases Control and Prevention (CDC) has also stated that it is not yet known whether non-respiratory body fluids from an infected person

<sup>1</sup> Reproduction Group, Department of Microbiology and Parasitology, Medical School, University of Antioquia, Antioquia, Colombia

<sup>2</sup> Department of Basic Medical Sciences, Mohammed Bin Rashid University of Medicine and Health Sciences, Dubai, UAE

<sup>3</sup> Division of Medical Physiology, Stellenbosch University, Tygerberg, South Africa

<sup>4</sup> Immunovirology Group, Department of Microbiology and Parasitology, Medical School, University of Antioquia, Antioquia, Colombia

## KEY WORDS

COVID-19  
Fertility  
SARS-CoV-2  
Sperm  
Virus

including vomit, urine, breast milk or semen can contain viable, infectious viruses (CDC, 2020). In Italy, one of the early COVID-19 epicenters, authorities are mandating that all gamete donors should be interviewed concerning the presence of respiratory symptoms and/or recent travel to high-risk areas, even in the absence of evidence of transmission of the virus within reproductive cells. A 2-week suspension from the end of symptoms is implemented in the case of donors showing respiratory symptoms, while a similar 2-week suspension is required from donors who have returned from an area at risk (La Marca et al., 2020).

Furthermore, based on current knowledge, it is of paramount importance that individuals who could potentially be infected take all possible care to mitigate the likely risk of passing on infection through sexual intercourse. Precautionary measurements should be practiced by candidates who have not yet been tested positive for SARS-CoV-2, and also by those previously infected, even after the signs and symptoms have resolved. It is recommended to avoid possible transfection by refraining from sexual relations without protection, as the virus could persist for a longer period of time in the male reproductive system compared with other body fluids, rendering the testis a potential viral reservoir.

## REFERENCES

Center for Diseases Control and Prevention (CDC), 2020. Clinical questions about COVID-19. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/faq.html>

Society for Assisted Reproductive Technology (SART), 2020. SART and ASRM issue advice for infertility patients concerning the novel coronavirus (COVID-19). <https://www.Sart.Org/news-and-publications/news-and-research/press-releases-and-bulletins/sart-and-asrm-issue-advice-for-infertility-patients-concerning-the-novel-coronavirus-covid-19>.

Cardona Maya, W.D., Du Plessis, S.S., Velilla, P.A. **Semen as virus reservoir?** JARG 2016; 33: 1255-1256

Cardona-Maya, W.D., Velilla Hernandez, P.A., Henao, D.E. **Male ebola survivors: Do not forget to use a condom!** Reprod. Sci. 2019; 26: 1326

Ding, Y., He, L., Zhang, Q., Huang, Z., Che, X., Hou, J., Wang, H., Shen, H., Qiu, L., Li, Z., Geng, J., Cai, J., Han, H., Li, X., Kang, W., Weng, D., Liang, P., Jiang, S. **Organ distribution of severe acute respiratory syndrome (SARS) associated coronavirus (SARS-CoV) in SARS patients: implications for pathogenesis and virus transmission pathways.** J. Pathol. 2004; 203: 622-630

Fan, C., Li, K., Ding, Y., Lu, L.W., Wang, J. **ACE2 expression in kidney and testis may cause kidney and testis damage after 2019-nCoV infection.** medRxiv. 2020. doi:10.1101/2020.02.12.20022418

Feldmann, H. **Virus in semen and the risk of sexual transmission.** N. Engl. J. Med. 2018; 378: 1440-1441

Gu, J., Gong, E., Zhang, B., Zheng, J., Gao, Z., Zhong, Y., Zou, W., Zhan, J., Wang, S., Xie, Z., Zhuang, H., Wu, B., Zhong, H., Shao, H., Fang, W., Gao, D., Pei, F., Li, X., He, Z., Xu, D., Shi, X., Anderson, V.M., Leong, A.S. **Multiple organ infection and the pathogenesis of SARS.** J. Exp. Med. 2005; 202: 415-424

La Marca, A., Niederberger, C., Pellicer, A., Nelson, S.M. **COVID-19: lessons from the Italian reproductive medical experience.** Fertil. Steril. 2020. doi:10.1016/j.fertnstert.2020.03.021

Rothe, C., Schunk, M., Sothmann, P., Bretzel, G., Froeschl, G., Wallrauch, C., Zimmer, T., Thiel, V., Janke, C. **Transmission of 2019-nCoV infection from an asymptomatic contact in Germany.** N. Engl. J. Med. 2020; 382: 970-971

Sigurðardóttir, Ó., Kolbjørnsen, O., Lutz, H. **Orchitis in a cat associated with coronavirus infection.** J. Comp. Pathol. 2001; 124: 219-222

Xu, J., Qi, Lihua, Chi, Xiaochun, Yang, Jingjing, Wei, Xiaohong, Gong, Encong, Peh, Suatcheng, Gu, Jiang **Orchitis: A complication of severe acute respiratory syndrome (SARS).** Biol. Reprod. 2006; 74: 410-416

Zhao, J.M., Zhou, G.D., Sun, Y.L., Wang, S.S., Yang, J.F., Meng, E.H., Pan, D., Li, W.S., Zhou, X.S., Wang, Y.D., Lu, J.Y., Li, N., Wang, D.W., Zhou, B.C., Zhang, T.H. **[Clinical pathology and pathogenesis of severe acute respiratory syndrome].** Zhonghua Shi Yan He Lin Chuang Bing Du Xue Za Zhi [Chin. J. Exp. and Clin. Vir.] 2003; 17: 217-221