

# Corrigendum: Variation in the *HLA-F* gene locus with functional impact is associated with pregnancy success and time-to-pregnancy after fertility treatment

Cæcilie H. Langkilde<sup>1,2,†</sup>, Line Lynge Nilsson<sup>1,2,‡</sup>, Nanna Jørgensen<sup>1,2</sup>, Tina Funck<sup>1,2</sup>, Trine L. Perin<sup>2,3</sup>, Micha B. Hornstrup<sup>2,3</sup>, Thomas Høst<sup>2,3,†</sup>, Thomas Scheike<sup>4</sup>, Anette Lindhard<sup>2,3</sup>, and Thomas Vauvert F. Hviid<sup>1,2,\*</sup>

<sup>1</sup>Centre for Immune Regulation and Reproductive Immunology (CIRRI), Department of Clinical Biochemistry, The ReproHealth Research Consortium ZUH, Zealand University Hospital, Roskilde, Denmark <sup>2</sup>Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark <sup>3</sup>The Fertility Clinic, Department of Obstetrics and Gynaecology, The ReproHealth Research Consortium ZUH, Zealand University Hospital, Køge, Denmark <sup>4</sup>Section of Biostatistics, Department of Public Health, University of Copenhagen, Copenhagen, Denmark

\*Correspondence address. Department of Clinical Biochemistry, Centre for Immune Regulation and Reproductive Immunology (CIRRI), The ReproHealth Research Consortium ZUH, Zealand University Hospital, and Department of Clinical Medicine, University of Copenhagen, 10 Sygehusvej, DK-4000 Roskilde, Denmark. Tel: +45-4732-5622; E-mail: tvh@regionsjaelland.dk

*Hum Reprod* 2020;**35**:705–717

The authors would like to apologise for errors in Table I of the above article.

In cycle 4, the number of IUI is 16 and not 17. The total number of women considered for analyses for the cycle in question is thus 50 instead of 51. Thereby, the total number of cycles included in the study was 387.

In cycle 3, 16 achieved pregnancy instead of 13; in cycle 5, 3 achieved pregnancy instead of 5; in cycle 6, 3 achieved pregnancy instead of 6;

in cycle 7, 3 achieved pregnancy instead of 5; in cycle 8, 2 achieved pregnancy instead of 3.

The correct numbers have been used for all the analyses and the authors would like to assure readers that changes made in Table I do not affect any results, conclusions or content of the article.

A corrected version of Table I is available below. The electronic version of this article has been updated at <https://doi.org/10.1093/humrep/dez276>.

<sup>†</sup>Deceased 11 June 2019.

<sup>‡</sup>These two authors contributed equally to this work.

**Table 1** Summary of ART cycles during the first 2 years of treatment.

Type of ART treatment:	Cycle 1 n (%)	Cycle 2 n (%)	Cycle 3 n (%)	Cycle 4 n (%)	Cycle 5 n (%)	Cycle 6 n (%)	Cycle 7 n (%)	Cycle 8 n (%)	Cycle 9 n (%)	Cycle 10 n (%)
IUI	86 (84.3)	67 (80.7)	47 (69.1)	16 (32.0)	8 (22.2)	4 (18.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
IVF	4 (3.9)	5 (6.0)	9 (13.2)	14 (28.0)	9 (25.0)	5 (22.7)	4 (26.7)	1 (14.3)	0 (0.0)	0 (0.0)
ICSI	11 (10.8)	8 (9.6)	8 (11.8)	16 (32.0)	15 (41.7)	7 (31.8)	9 (60.0)	3 (42.9)	1 (33.3)	1 (100.0)
Frozen embryo replacement IVF	0 (0.0)	0 (0.0)	1 (1.5)	3 (6.0)	2 (5.6)	1 (4.5)	0 (0.0)	1 (14.3)	0 (0.0)	0 (0.0)
Frozen embryo replacement ICSI	0 (0.0)	3 (3.6)	2 (2.9)	1 (2.0)	2 (5.6)	4 (18.2)	1 (6.7)	2 (28.6)	2 (66.7)	0 (0.0)
Spontaneously achieved pregnancy within 2 years with a live birth and after end of treatment <sup>a)</sup>	1 (1.0)	0 (0.0)	1 (1.5)	0 (0.0)	0 (0.0)	1 (4.5)	1 (6.7)	0 (0.0)	0 (0.0)	0 (0.0)
<b>Total number of women considered for analyses for the cycle in question<sup>b)</sup></b>	102 (100)	83 (100)	68 (100)	50 (100)	36 (100)	22 (100)	15 (100)	7 (100)	3 (100)	1 (100)
<b>Total number of couples that achieved pregnancy with a live birth after treatment within 2 years</b>	15	10	16	4	3	3	3	2	1	0

<sup>a)</sup> A pregnancy with a live birth achieved spontaneously within the first 2 years of treatment is considered as one treatment cycle. Regarding the one pregnancy in cycle 1, the couple was enrolled for ART treatment but conceived spontaneously before start of treatment.

<sup>b)</sup> The total number of cycles included in the study was 387.