

Birthweight and other perinatal outcomes of singletons conceived after assisted reproduction compared to natural conceived singletons in couples with unexplained subfertility: follow-up of two randomized clinical trials

J.A. Wessel¹, F. Mol¹*, N.A. Danhof¹, A.J. Bensdorp¹,
R.I. Tjon-Kon Fat¹, F.J.M. Broekmans², A. Hoek³, B.W.J. Mol⁴,
M.H. Mochtar¹, and M. van Wely¹; INeS and SUPER Study Group[†]

¹Centre for Reproductive Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam 1105 AZ, the Netherlands ²Centre for Reproductive Medicine, University Medical Centre Utrecht, Utrecht 3508 GA, the Netherlands ³Department of Obstetrics and Gynaecology, University of Groningen, University Medical Centre Groningen, Groningen 9713 GZ, the Netherlands ⁴Department of Obstetrics and Gynaecology, Monash University, Monash Medical Centre, Clayton, VIC 3168, Australia

*Correspondence address. Centre for Reproductive Medicine, Amsterdam UMC, University of Amsterdam, Meibergdreef 9, Amsterdam 1105 AZ, the Netherlands. E-mail: f.mol@amsterdamumc.nl

Submitted on July 2, 2020; resubmitted on October 5, 2020; editorial decision on October 12, 2020

STUDY QUESTION: Does assisted reproduction, such as ovarian stimulation and/or laboratory procedures, have impact on perinatal outcomes of singleton live births compared to natural conception in couples with unexplained subfertility?

SUMMARY ANSWER: Compared to natural conception, singletons born after intrauterine insemination with ovarian stimulation (IUI-OS) had a lower birthweight, while singletons born after IVF had comparable birthweights, in couples with unexplained subfertility.

WHAT IS KNOWN ALREADY: Singletons conceived by assisted reproduction have different perinatal outcomes such as low birthweight and a higher risk of premature birth than naturally conceived singletons. This might be due to the assisted reproduction, such as laboratory procedures or the ovarian stimulation, or to an intrinsic factor in couples with subfertility.

STUDY DESIGN, SIZE, DURATION: We performed a prospective cohort study using the follow-up data of two randomized clinical trials performed in couples with unexplained subfertility. We evaluated perinatal outcomes of 472 live birth singletons conceived after assisted reproduction or after natural conception within the time horizon of the studies.

PARTICIPANTS/MATERIALS, SETTING, METHODS: To assess the possible impact of ovarian stimulation we compared the singletons conceived after IUI with FSH or clomiphene citrate (CC) and IVF in a modified natural cycle (IVF-MNC) or standard IVF with single embryo transfer (IVF-SET) to naturally conceived singletons in the same cohorts. To further look into the possible effect of the laboratory procedures, we put both IUI and IVF groups together into IUI-OS and IVF and compared both to singletons born after natural conception. We only included singletons conceived after fresh embryo transfers. The main outcome was birthweight presented as absolute weight in grams and gestational age- and gender-adjusted percentiles. We calculated differences in birthweight using regression analyses adjusted for maternal age, BMI, smoking, parity, duration of subfertility and child gender.

MAIN RESULTS AND THE ROLE OF CHANCE: In total, there were 472 live birth singletons. Of the 472 singleton pregnancies, 209 were conceived after IUI-OS (136 with FSH and 73 with CC as ovarian stimulation), 138 after IVF (50 after IVF-MNC and 88 after IVF-SET) and 125 were conceived naturally.

[†]The members of the INeS and SUPER Study Group are listed in the Appendix.

© The Author(s) 2020. Published by Oxford University Press on behalf of European Society of Human Reproduction and Embryology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Singletons conceived following IUI-FSH and IUI-CC both had lower birthweights compared to naturally conceived singletons (adjusted difference IUI-FSH –156.3 g, 95% CI –287.9 to –24.7; IUI-CC –160.3 g, 95% CI –316.7 to –3.8). When we compared IVF-MNC and IVF-SET to naturally conceived singletons, no significant difference was found (adjusted difference IVF-MNC 75.8 g, 95% CI –102.0 to 253.7; IVF-SET –10.6 g, 95% CI –159.2 to 138.1). The mean birthweight percentile was only significantly lower in the IUI-FSH group (–7.0 percentile, 95% CI –13.9 to –0.2). The IUI-CC and IVF-SET group had a lower mean percentile and the IVF-MNC group a higher mean percentile, but these groups were not significantly different compared to the naturally conceived group (IUI-CC –5.1 percentile, 95% CI –13.3 to 3.0; IVF-MNC 4.4 percentile, 95% CI –4.9 to 13.6; IVF-SET –1.3 percentile, 95% CI –9.1 to 6.4).

Looking at the laboratory process that took place, singletons conceived following IUI-OS had lower birthweights than naturally conceived singletons (adjusted difference –157.7 g, 95% CI –277.4 to –38.0). The IVF group had comparable birthweights with the naturally conceived group (adjusted difference 20.9 g, 95% CI –110.8 to 152.6). The mean birthweight percentile was significantly lower in the IUI-OS group compared to the natural group (–6.4 percentile, 95% CI –12.6 to –0.1). The IVF group was comparable (0.7 percentile, 95% CI –6.1 to 7.6).

LIMITATIONS, REASONS FOR CAUTION: The results are limited by the number of cases. The data were collected prospectively alongside the randomized controlled trials, but analyzed as treated.

WIDER IMPLICATIONS OF THE FINDINGS: Our data suggest IUI in a stimulated cycle may have a negative impact on the birthweight of the child and possibly on pre-eclampsia. Further research should look into the effect of different methods of ovarian stimulation on placenta pathology and pre-eclampsia in couples with unexplained subfertility using naturally conceived singletons in the unexplained population as a reference.

STUDY FUNDING/COMPETING INTEREST(S): Both initial trials were supported by a grant from ZonMW, the Dutch Organization for Health Research and Development (INeS 120620027, SUPER 80-83600-98-10192). The INeS study also had a grant from Zorgverzekeraars Nederland, the Dutch association of healthcare insurers (09-003). B.W.J.M. is supported by an NHMRC investigator Grant (GNT1176437) and reports consultancy for ObsEva, Merck Merck KGaA, Guerbet and iGenomix, outside the submitted work. A.H. reports grants from Ferring Pharmaceutical company (the Netherlands), outside the submitted work. F.J.M.B. receives monetary compensation as a member of the external advisory board for Merck Serono (the Netherlands), Ferring Pharmaceuticals BV (the Netherlands) and Gedeon Richter (Belgium), he receives personal fees from educational activities for Ferring BV (the Netherlands) and for advisory and consultancy work for Roche and he receives research support grants from Merck Serono and Ferring Pharmaceuticals BV, outside the submitted work. The remaining authors have nothing to disclose.

TRIAL REGISTRATION NUMBER: INeS study Trial NL915 (NTR939); SUPER Trial NL3895 (NTR4057)

Key words: unexplained subfertility / ovarian stimulation / laboratory procedures / perinatal outcomes / birthweight

Introduction

Up to a quarter of all couples who are unable to conceive after 12 months of unprotected intercourse, and where routine fertility investigations show no abnormalities are diagnosed with unexplained subfertility (Tjon-Kon-Fat *et al.*, 2016). These couples are treated with either intrauterine insemination with ovarian stimulation (IUI-OS) or IVF. The 18th ESHRE report on assisted reproduction shows that in the year 2014 a total of 120 789 IUI cycles and 146 148 IVF cycles were performed in 39 European countries (De Geyter *et al.*, 2018), so both treatments are widely used. The prediction model of Hunault can be used in these couples with unexplained subfertility to differentiate between couples that have favorable chances of conceiving naturally and those that do not. The model is an externally validated synthesis model and includes female age, duration of subfertility, sperm motility, a previous pregnancy and referral status to contribute in the shared decision-making process when choosing the best treatment option. In couples with the 1-year prognosis to conceive naturally below 30%, assisted reproduction may improve efficacy but solid evidence is missing (Hunault *et al.*, 2004; Wang *et al.*, 2019). Notably, many couples still conceive naturally in between treatment cycles (van Eekelen *et al.*, 2018). To decide which strategy is preferred; i.e. either continue with expectant management or start with IUI or proceed directly to IVF, safety aspects should also be taken into account.

Initial concerns about the safety of IVF were mainly attributed to a higher rate of multiple births (Pinborg *et al.*, 2004). The trend toward single embryo transfer (SET) to prevent multiple pregnancies decreased the number of perinatal complications from assisted reproduction (Kallen *et al.*, 2010). Yet, large observational studies have shown that singletons born after different types of assisted reproduction still have different perinatal outcomes such as lower birthweight (LBW) and a higher risk of being born premature compared to naturally conceived singletons (Pinborg *et al.*, 2013; Dunietz *et al.*, 2015). As these studies usually compared perinatal outcomes of singletons in subfertile couples to fertile couples, it remains difficult to determine whether these perinatal outcomes are assisted reproduction procedures related, or the result of a possible intrinsic factor in the subfertile couples (Pandey *et al.*, 2012; Ombelet *et al.*, 2016).

Several causes of the different perinatal outcomes of singletons following assisted reproduction have been put forward. First, the ovarian stimulation could be a potential risk factor for abnormal placentation leading to lower birthweights (Pereira *et al.*, 2015). Second, the laboratory procedures of assisted reproduction appear to increase the risks (Lambert, 2003). Third, subfertile couples may have an intrinsic higher risk of adverse perinatal outcomes (Pelinck *et al.*, 2010; Messerlian *et al.*, 2013).

To overcome the intrinsic effect of subfertility, the outcomes after assisted reproduction and naturally conceived children should be

evaluated within the unexplained subfertile population. As we know of this comparison has not been reported yet and the current study could be a contribution to the scientific literature. We evaluated in a population of couples with unexplained subfertility if ovarian stimulation and/or artificial laboratory procedures affected perinatal outcomes of the live born singletons compared to natural conception in the same cohort, therefore leaving the intrinsic factor of subfertile couples out of the equation.

Materials and methods

Study design

We performed a prospective cohort study using the follow-up data of two randomized clinical trials (RCT) in couples with unexplained subfertility and mild male subfertility (INeS trial NTR 939, SUPER study NTR 4057). Both the initial trials were approved by the medical Ethical Committee of the Academic Medical Centre and the Central Committee on Research involving Human Subjects (CCMO) (INeS: NL12782.018.07; Super: NL 43131-018-13). Participants provided written informed consent for use of the follow-up data during entry in both the RCTs.

The INeS trial was a Dutch multicenter randomized controlled non-inferiority trial, they included 602 couples and compared the effectiveness of IUI-OS, modified natural cycle IVF (IVF with monofollicular growth using a GnRH antagonist and gonadotropins when follicle dominance occurred) (IVF-MNC), and standard IVF with controlled hyperstimulation and SET (IVF-SET) (Bensdorp *et al.*, 2015). The SUPER study was a Dutch multicenter randomized controlled superiority trial, they included 738 women and compared the effectiveness of IUI with FSH versus clomiphene citrate (CC) (Danhof *et al.*, 2018). The primary outcomes were healthy singletons (INeS trial) and ongoing pregnancies (SUPER study).

Study population

Both RCTs included couples diagnosed with unexplained or mild male subfertility. Unexplained subfertility was defined as a failure to conceive after 1 year of regular unprotected intercourse where fertility investigations showed an ovulatory cycle and at least one patent fallopian tube. Mild male subfertility was described as a prewash total motile sperm count of at least 3 million. Only couples with an unfavorable prognosis, defined as a 12-month prognosis at natural conception according to the model of Hunault lower than 30%, were included (Hunault *et al.*, 2004). Both trials excluded women with anovulation, double-sided tubal disease, polycystic ovarian syndrome and other endocrine diseases at baseline (such as Cushing's syndrome or diabetes mellitus type I). The INeS study also excluded women with severe endometriosis, premature ovarian failure and women older than 38 years. The SUPER trial excluded women older than 43 years. We only included the singletons conceived after fresh embryo transfers.

Outcomes

The main outcome is birthweight presented as absolute weight in grams and gestational age- and gender-adjusted percentiles. Other outcomes were LBW (birthweight <2500 g), very LBW (birthweight

<1500 g), small for gestational age (SGA) (birthweight <10th and <3rd of the national reference), high birthweight (birthweight >4000 g), large for gestation age (birthweight >90th centile of the national reference), preterm birth (delivery <37, <32 and <28 weeks of gestation) split up in iatrogenic or spontaneous, congenital anomalies and pre-eclampsia.

We assessed the possible impact of ovarian stimulation by comparing the singletons conceived after IUI with FSH or CC and IVF-MNC or IVF-SET to naturally conceived singletons in the same cohorts. To further look into the possible effect of the laboratory procedures, we put both IUI and IVF groups together into IUI-OS and IVF. Then we compared both groups to singletons born after natural conception during the study period.

To compare the birthweights in our study with the birthweights of singleton live births from the general Dutch population, we calculated gestational age- and gender-adjusted percentiles based on the Dutch reference curves. These birthweight percentiles allow us to evaluate birth weight independent of gestational age and gender. The Dutch reference curves are based on a low-risk population (infants with one or more risk factors for abnormal fetal growth were excluded) and included 1 629 776 live birth singleton infants (2000–2014) (Hofstiezer *et al.*, 2019).

Statistical analysis

Continuous variables were expressed as mean \pm standard deviation and categorical variables were expressed as number and percentage. We performed our primary analysis on basis of the as-treated population, grouped as IUI-FSH, IUI-CC, IVF-MNC, IVF-SET and natural conception. (The intention to treat numbers as retrieved from the INeS and SUPER trial are available in [Supplementary Tables SI and SII](#).) For all analyses, the naturally conceived singletons were the control group.

We assessed the impact of ovarian stimulation and laboratory procedures on the birthweight of a singleton, by using linear regression analyses with adjustment for maternal age, pre-conception BMI, smoking, parity, subfertility length and child gender. The decision to enter these factors was made a priori based on the known influence on birthweight. The binary outcomes were compared using Fisher Exact. *P*-values below 0.05 are considered as statistically significant differences. We used SPSS (version 25.0) for all statistical analyses.

Results

In total, both trials reported 541 ongoing pregnancies. We excluded 36 multiple gestations (12 in the IVF group and 23 in the IUI-OS group). Twenty-five women with an ongoing singleton pregnancy were lost to follow-up. Eight women had no live birth between 17 and 37 weeks of gestation, no further information is available. Of the remaining 472 live birth singletons, 136 were conceived after IUI-FSH, 73 after IUI-CC, 50 after IVF-MNC, 88 after IVF-SET and 125 were conceived naturally. We summarized the study sample in a flowchart (Fig. 1). The baseline characteristics are depicted in [Table I](#). The IVF-SET group differs significantly in pre-conception BMI and duration of subfertility, the IUI-FSH group was more primary subfertile compared to the naturally conceived group.




Figure 1. Study flowchart. IUI-CC, intrauterine insemination with clomiphene citrate; IUI-FSH, intrauterine insemination with follicle stimulating hormone; IUI-OS, intrauterine insemination with ovarian stimulation; IVF-MNC, IVF in a modified natural cycle; IVF-SET, standard IVF with single embryo transfer.

Birthweight

First we compared the birthweights in the different ovarian stimulation groups to natural conception adjusted for age, BMI, smoking, parity, subfertility length and child gender. The mean birthweights were lower in both the IUI-FSH and IUI-CC groups when compared to the naturally conceived group, after adjustment, the difference remained significant (adjusted difference IUI-FSH –156.3 g, 95% CI –287.9 to –24.7; IUI-CC –160.3 g, 95% CI –316.7 to –3.8). The mean birthweights following IVF-MNC and IVF-SET were not significantly different compared to the naturally conceived group (adjusted difference IVF-MNC 75.8 g, 95% CI –102.0 to 253.7; IVF-SET –10.6 g, 95% CI –159.2 to 138.1) (Table II).

We subsequently compared the birthweights of the groups based on the laboratory procedures that took place. The birthweight of singletons born after IUI-OS was lower than in the naturally conceived group (adjusted difference –157.7 g, 95% CI –277.4 to –38.0). The mean birthweight in the IVF group was comparable to the naturally conceived group (adjusted difference 20.9 g, 95% CI –110.8 to 152.6) (Table III).

> Percentiles

We compared the birthweights of our singletons to singletons in the low-risk Dutch population by calculating the gestational age- and gender-adjusted percentiles. We found a significant lower birthweight percentile following IUI-FSH compared to natural conception (mean difference –7.0 percentile, 95% CI –13.9 to –0.2) while in the IUI-CC group the difference was not significant (mean difference –5.1 percentile, 95% CI –13.3 to 3.1). The birthweight percentiles for

singletons born after IVF-MNC and IVF-SET were comparable with singletons born after natural conception, which is almost average (Tables II and III, Supplementary Figs S1 and S2).

Other outcomes

With respect to ovarian stimulation, the IUI-FSH group had significantly more mothers with pre-eclampsia during their pregnancy compared to the naturally conceived group (8.6% vs. 2.6% $P=0.05$). Six of the 17 mothers who suffered from pre-eclampsia had an SGA singleton. In total, 46 singletons were conceived with donor semen. One of these mothers developed pre-eclampsia, she conceived after IUI-FSH and had a singleton with a normal birthweight.

The IUI-CC group had the most singletons with very LBW (5.5% vs. 0.8% $P=0.04$) compared to naturally conceived singletons. Of the singletons conceived after IUI-FSH and IUI-CC a significantly smaller number were born with very high birthweight compared to natural conception (6.7% and 5.5% vs. 15.2%, $P=0.03$ and $P=0.04$) (Table II).

When comparing the laboratory procedures, the IUI procedure resulted in less very high birthweight (>4000 g) when compared to natural (6.3% vs. 15.2%, $P=0.01$). We found no differences between IVF and natural conception (Table III).

Twenty-eight women had a preterm delivery before 37 weeks. In 16 women (57.1%), the onset of premature labor was spontaneous and started with contractions ($n=10$) or preterm premature rupture of membranes PPROM[AQ: Please spell out PPROM (if necessary).] followed by natural contractions ($n=6$). We considered the other

Table I Baseline characteristics of couples based on the mode of reproduction (n = 472).

	IUI-FSH (n = 136)	IUI-CC (n = 73)	IVF-MNC (n = 50)	IVF-SET (n = 88)	Natural (n = 125)
Age in years	32.67 (3.5)	32.82 (3.4)	33.51 (2.9)	32.8 (3.4)	32.75 (3.6)
BMI in kg/m ² ^a	24.7 (4.2)	24.4 (3.8)	23.1 (3.3)	23.0 (3.3) [*]	24.0 (4.2)
Duration subfertility in months ^b	34.7 (16.0)	28.1 (11.8)	30.8 (9.2)	36.6 (15.8) [*]	31.8 (14.2)
Smoking					
Yes	26 (19.1)	9 (12.3)	6 (12.0)	22 (20.2)	23 (18.4)
No	110 (80.9)	64 (87.7)	44 (88.0)	87 (78.8)	102 (81.6)
Primary subfertility					
Yes	120 (88.2) ⁺	59 (80.8)	39 (78.0)	86 (78.9)	93 (74.4)
No	16 (11.8)	14 (19.2)	11 (22.0)	23 (21.1)	32 (25.6)
Total motile sperm count ($\times 10^6$)	72.1 (69.0)	84.0 (113.8)	86.6 (97.1)	79.5 (95.2)	73.0 (72.2)

Data are n (%) or mean (SD).

^aLinear regression analyses P < 0.05 compared to natural.

⁺Chi-square test P < 0.05 compared to natural.

^an = 459.

^bn = 469.

IUI-CC, intrauterine insemination with clomiphene citrate; IUI-FSH, intrauterine insemination with FSH; IVF-MNC, IVF in a modified natural cycle; IVF-SET, standard IVF with single embryo transfer.

premature births (n = 12/28; 42.9%) as 'iatrogenic' since the onset was not spontaneous but the result of a decision by the clinician on various indications (i.e. indicated primary cesarean section (n = 7) or indicated induction of labor (n = 5)). There were no women with PPROM in the iatrogenic group. The distribution with details on preterm birth over the different treatment groups can be found in Tables II and III.

Congenital anomalies

There were five cases of congenital anomalies that we classified according to the ICD-10 classification (international classification of diseases, 10th revision). There were two cases of esophageal atresia (Q39) in the IUI-FSH group. One case of Aicardi syndrome (Q4) occurred in the IUI-CC group. In the IVF-MNC group, one case of Moebius syndrome (Q87) occurred and in the IVF-SET group, one case of Down syndrome (Q90) occurred. None of the naturally conceived singletons had congenital anomalies. We performed a linear regression analyses with adjustment for congenital anomalies. The mean difference in birthweight remained significantly lower in the IUI-FSH and IUI-CC groups compared to naturally conceived singletons (adjusted difference IUI-FSH – 153.4 g, 95% CI –281.0 to –25.9; IUI-CC – 166.8 g, 95% CI –318.4 to –15.2) and in the IVF-MNC and IVF-SET groups the mean birthweights were comparable to naturally conceived singletons (adjusted difference IVF-MNC 67.0 g, 95% CI –105.2 to 239.3; IVF-SET – 11.3 g, 95% CI –154.4 to 131.8).

Discussion

We evaluated whether ovarian stimulation or laboratory procedures in assisted reproduction affected the perinatal outcomes of singleton live births in couples with unexplained subfertility. Singletons conceived following IUI-OS had between 38 and 280 g lower birthweights than

naturally conceived singletons. The observed birthweight difference was irrespective of whether FSH or CC was used. The difference in birthweight was reflected by more very low birthweight (<1500 g) and less very high birthweight (>4000 g) in singletons conceived by IUI-OS than in natural conceptions. Singletons conceived by IVF, whether IVF-MNC or IVF-SET, did not have lower birthweights compared to naturally conceived singletons in unexplained subfertile couples. These results were confirmed when calculating the birthweight percentile relative to the low-risk Dutch reference population of singletons.

Based on literature, we expected to find lower birthweights in both the IUI-OS and IVF group compared to natural conception (Bentzen *et al.*, 2019). However, these studies compared children conceived after assisted reproduction with children conceived after natural conception from fertile populations. Studies which did look at subfertile couples did not only include couples with unexplained subfertility (Pelinck *et al.*, 2010). This may explain why we found no indication for a lower birthweight following IVF. Because, in contrast to most studies, we excluded couples with other causes of subfertility such as polycystic ovary syndrome or other endocrine disorders which maybe influence birthweight (Messerlian *et al.*, 2013). This theory is supported by data from Isaksson *et al.* (2002), a case control study where the perinatal outcomes of singletons born in couples with unexplained subfertility who conceived after IVF/ICSI were compared to perinatal outcomes of singletons born in couples without subfertility who conceived naturally. There was no difference in mean birthweight between both groups in this study. The overall perinatal outcomes of singletons among couples with unexplained subfertility treated with IVF/ICSI were similar to the perinatal outcomes of singletons born after natural conception in couples without fertility problems (Isaksson *et al.*, 2002).

The strengths of our study are that firstly we only used live birth singletons in our comparisons and secondly the studied population was homogenous in that all the included couples shared the same profile and suffered from unexplained subfertility with poor chances on

Table II Outcomes of singletons based on the mode of reproduction and the type of ovarian stimulation of the mother (n = 472).

	IUI-FSH (n = 136)	IUI-CC (n = 73)	IVF-MNC (n = 50)	IVF-SET (n = 88)	Natural (n = 125)
Mean birthweight in grams (SD)	3296 (492)	3283 (667)	3515 (465)	3439 (418)	3453 (548)
Mean birthweight difference compared to natural (95% CI)	–157.1 (–284.5 to –29.8)*	–170.2 (–321.7 to –18.8)*	62.0 (–110.0 to 234.0)	–14.2 (–157.2 to 128.9.)	–
Adjusted mean birthweight difference compared to natural (95% CI)	–156.3 (–287.9 to –24.7)*	–160.3 (–316.7 to –3.8)*	75.8 (–102.0 to 253.7)	–10.6 (–159.2 to 138.1)	–
Mean birthweight percentile (SD) ^a	42.4 (27.4)	44.3 (29.9)	53.8 (28.0)	48.1 (27.8)	49.4 (28.5)
Mean percentile difference compared to natural (95% CI)	–7.0 (–13.9 to –0.2)*	–5.1 (–13.3 to 3.0)	4.4 (–4.9 to 13.6)	–1.3 (–9.1 to 6.4)	–
Mean gestational age in weeks (SD)	39.3 (2.0)	39.1 (2.5)	39.3 (1.4)	39.2 (1.4)	39.4 (2.0)
	n (%)	n (%)	n (%)	n (%)	n (%)
Infant sex					
Male	59 (43.4)	40 (54.8)	19 (38.0)	45 (51.1)	64 (51.2)
Female	77 (56.6)	33 (45.2)	31 (62.0)	43 (48.9)	61 (48.8)
SGA (<p10)	17 (12.6)	10 (13.7)	4 (8.0)	6 (6.8)	15 (12.0)
SGA (<p3)	4 (3.0)	5 (6.8)	1 (2.0)	1 (1.1)	3 (2.4)
Low birthweight (<2500 g)	5 (3.7)	7 (9.6)	1 (2.0)	2 (2.3)	7 (5.6)
Very low birthweight (<1500 g)	2 (1.5)	4 (5.5) ⁺	0 (0.0)	0 (0.0)	1 (0.8)
Preterm birth (<37 w)	9 (6.6)	8 (11.0)	2 (4.0)	3 (3.4)	6 (4.8)
<i> iatrogenic</i>	4 (44.4)	3 (37.5)	2 (100.0)	1 (33.3)	2 (33.3)
<i> Spontaneous</i>	5 (55.6)	5 (62.5)	0 (0.0)	2 (66.7)	4 (66.7)
Preterm birth (<32 w)	2 (1.5)	3 (4.1)	0 (0.0)	0 (0.0)	2 (1.6)
<i> iatrogenic</i>	0 (0.0)	2 (66.7)	0 (0.0)	1 (50.0)	1 (50.0)
<i> Spontaneous</i>	2 (100.0)	1 (33.3)	0 (0.0)	1 (50.0)	1 (50.0)
Preterm birth (<28 w)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.8)
<i> iatrogenic</i>				1 (100.0)	
<i> Spontaneous</i>				0 (0.0)	
High birthweight (>4000 g)	9 (6.7) ⁺	4 (5.5) ⁺	6 (12.0)	7 (8.0)	19 (15.2)
LGA (>p90)	5 (3.7)	5 (6.8)	5 (10.0)	8 (9.1)	9 (6.9)
Congenital anomalies	2 (1.5)	1 (1.4)	1 (2.0)	1 (1.1)	0 (0.0)
Pre-eclampsia ^b	10 (8.6) ^{c,+}	1 (1.6)	2 (4.0)	1 (1.1)	3 (2.6)

^aGestational age- and gender-adjusted percentile based on the Dutch reference curve (Hofstiezer et al., 2019).^bn = 452.^cOne woman conceived with donor semen.^{*}Linear regression analyses P < 0.05 compared to natural.[†]Chi-square test P < 0.05 compared to natural.

conceiving naturally within 12 months after fertility work-up according to the model of Hunault (Hunault score <30%). A limitation of this study is that the power of this study is relatively low due to the rather small numbers of cases, so the results have to be interpreted with caution. A second limitation of this study is that we were not able to adjust for other confounding factors such as high blood pressure, gestational diabetes mellitus or paternal characteristics because these variables were not selected for collection in the case record forms of the original trials. Nevertheless, we found no evidence in our study to suggest a clinically important reduction in birthweight due to IVF in the population of unexplained subfertility. Whether the FSH or CC or the

IUI process truly has a negative impact on birthweight needs to be investigated further.

We observed a relatively high number of women who developed pre-eclampsia in the IUI-FSH group. This might be a chance finding as the expected incidence of pre-eclampsia is low and the number of singletons in our study was limited. In a previous retrospective cohort study, there was no evidence for an association between IUI and pre-eclampsia in a general subfertile population (Chen et al., 2009). Women conceiving with donor semen might have a higher risk of pre-eclampsia compared to partners semen (Gonzalez-Comadran et al., 2014). In our study, we did not find this.

Table III Outcomes of singletons based on the mode of reproduction and laboratory procedures that took place (n = 472).

	IUI-OS (n = 209)	IVF (n = 138)	Natural (n = 125)
Mean birthweight in grams (SD)	3291 (558)	3467 (436)	3453 (548)
Mean birthweight difference compared to natural (95% CI)	-161.7 (-277.8 to -45.6)*	13.4 (-113.3 to 140.2)	-
Adjusted mean birthweight difference compared to natural (95% CI)	-157.7 (-277.4 to -38.0)*	20.9 (-110.8 to 152.6)	-
Mean birthweight percentile ^a (SD)	43.1 (28.2)	50.2 (27.9)	49.4 (28.5)
Mean percentile difference compared to natural (95% CI)	-6.4 (-12.6 to -0.1)*	0.7 (-6.1 to 7.6)	-
Mean gestational age in weeks (SD)	39.2 (2.2)	39.3 (1.4)	39.4 (2.0)
	n (%)	n (%)	n (%)
Infant sex			
Male	99 (47.4)	64 (46.4)	64 (51.2)
Female	110 (52.6)	74 (53.6)	61 (48.8)
SGA (<p10)	27 (13.0)	10 (7.2)	15 (12.0)
SGA (<p3)	9 (4.3)	2 (1.4)	3 (2.4)
Low birthweight (<2500 g)	12 (5.8)	3 (2.2)	7 (5.6)
Very low birthweight (<1500 g)	6 (2.9)	0 (0.0)	1 (0.8)
Preterm birth (<37 w)	17 (8.6)	5 (3.6)	6 (4.8)
<i>iatrogenic</i>	7 (41.2)	3 (60.0)	2 (33.3)
<i>Spontaneous</i>	10 (58.8)	2 (40.0)	4 (66.7)
Preterm birth (<32 w)	5 (2.4)	0 (0.0)	2 (1.6)
<i>iatrogenic</i>	2 (40.0)		1 (50.0)
<i>Spontaneous</i>	3 (60.0)		1 (50.0)
Preterm birth (<28 w)	0 (0.0)	0 (0.0)	1 (0.8)
<i>iatrogenic</i>			1 (100.0)
<i>Spontaneous</i>			0 (0.0)
High birthweight (>4000 g)	13 (6.3) ⁺	13 (9.4)	19 (15.2)
LGA (>p90)	10 (4.8)	13 (9.4)	9 (7.2)
Congenital anomalies	3 (1.4)	2 (1.4)	0 (0.0)
Pre-eclampsia ^b	11 (6.2) ^c	3 (2.2)	3 (2.6)

^aGestational age- and gender-adjusted percentile based on the Dutch reference curve (Hofstiezer *et al.*, 2019).

^bn = 452.

^cOne woman conceived with donor semen.

*Linear regression analyses P < 0.05 compared to natural.

⁺Chi-square test P < 0.05 compared to natural.

IUI-OS, intrauterine insemination with ovarian stimulation; IVF, standard IVF with single embryo transfer; LGA, large for gestational age; SGA, small for gestational age.

It is unclear by what mechanism ovarian stimulation might influence birthweight. The general theory is that high estradiol concentrations can play a role, which can cause a negative effect on angiogenesis and placenta (Pereira *et al.*, 2015). We do not have data of the estradiol levels during stimulation, based on literature, we expect the estradiol levels to be higher in the IVF-SET group because this corresponds to the number of follicles which we expect to be highest in the IVF-SET group. This means that estradiol levels are probably not an explanation for the lower birthweights in the IUI-OS group (Joo *et al.*, 2010).

The only laboratory process that takes place during IUI treatments is processing semen. To our knowledge, there is no literature available that studied the influence of processed semen on perinatal outcomes

of singletons. Processing of semen also takes place at IVF and since we have not found a different mean birthweight in the IVF group compared to naturally conceived singletons we believe that processing semen and the *in-vitro* procedure do not affect the perinatal outcomes of singletons.

A possible explanation for the lower birthweight in the IUI-OS group might be vanishing twins. The remaining fetus after a vanishing twin has an increased risk for a lower birthweight and being born premature (Magnus *et al.*, 2017). In the absence of viable twin pregnancies, the development of multiple follicles in IUI-OS might still have resulted in vanishing twins, while vanishing twins are not to be expected following IVF-SET.

Conclusion

In conclusion, in singletons of couples with unexplained subfertility, we found lower mean birthweights when conception occurred after IUI-OS (irrespective of FSH or CC was used as ovarian stimulation) compared to natural conception. The singletons also had a lower mean birthweight than the singletons of the low-risk Dutch population. Our study suggests that in couples with unexplained subfertility, IVF results in comparable birthweights and other perinatal outcomes of singletons when compared to natural conception. Further large prospective cohort studies in an unexplained subfertile population are required to substantiate these findings.

Supplementary data

Supplementary data are available at *Human Reproduction* online.

Data availability

The data underlying this article will be shared on reasonable request to the corresponding author.

Acknowledgments

We thank all the couples, the hospitals and their staff that participated in the trials. For detailed information of participating hospitals see literature of the INeS trial and SUPER study.

Author's roles

J.A.W. is responsible for performing the analysis and drafted the paper. M.v.W. helped designing both initial trials. MM helped designing the super study. F.M. and M.v.W. supervised the analyses. All authors acquired the data from the participating centers, provided critical discussion, contributed in the preparation of the manuscript and approved the final version.

Funding

Both initial trials were supported by a grant from ZonMW, the Dutch Organization for Health Research and Development (INeS I20620027, SUPER 80-83600-98-10192). The INeS also had a grant from Zorgverzekerlaars Nederland, the Dutch association of healthcare insurers (09-003).

Conflict of interest

B.W.J.M. is supported by an NHMRC investigator Grant (GNT1176437) and reports consultancy for ObsEva, Merck Merck KGaA, Guerbet and iGenomix, outside the submitted work. A.H. reports grants from Ferring Pharmaceutical company (the Netherlands), outside the submitted work. F.J.M.B. receives monetary compensation as a member of the external advisory board for Merck Serono (the Netherlands), Ferring Pharmaceuticals BV (the Netherlands) and Gedeon Richter (Belgium), he receives personal fees from

educational activities for Ferring BV (the Netherlands) and for advisory and consultancy work for Roche and he receives research support grants from Merck Serono and Ferring Pharmaceuticals BV, outside the submitted work. The remaining authors have nothing to disclose.

Appendix

INeS study group members: A.J. Bensdorp, R.I. Tjon-Kon-Fat, C.A.M. Koks, G.J.E. Oosterhuis, A. Hoek, M.O. Verhoeven, F.J.M. Broekmans, H.R. Verhoeve, J.P. de Bruin, R. van Golde, B.J. Cohlen, M.D.A. Lambers, P.F. van Bommel, E. Slappendel, D.A.M. Perquin, J.M. Smeenk, M.J. Pelinck, J. Gianotten, D.A. Hoozemans, J.W.M. Maas, B.W.J. Mol and M van Wely.

SUPER study group members: N.A. Danhof, M. van Wely, C.A.M. Koks, H.R. Verhoeve, J.P. de Bruin, M.F.G. Verberg, M.H.A. van Hooff, B.J. Cohlen, C.F. van Heteren, K. Fleischer, J. Gianotten, J. van Disseldorp, J. Visser, F.J.M. Broekmans, B.W.J. Mol, D.P. van der Ham, N. Klijn, C.A.H. Janssen, J.M. van Rijn-van Weert, M. Twisk, M.J. Pelinck, D.A.M. Perquin, D.E.S. Boks, A. Sluijmer and M.H. Mochtar.

References

Bensdorp AJ, Tjon-Kon-Fat RI, Bossuyt PM, Koks CA, Oosterhuis GJ, Hoek A, Hompes PG, Broekmans FJ, Verhoeve HR, de Bruin JP et al. Prevention of multiple pregnancies in couples with unexplained or mild male subfertility: randomised controlled trial of in vitro fertilisation with single embryo transfer or in vitro fertilisation in modified natural cycle compared with intrauterine insemination with controlled ovarian hyperstimulation. *BMJ* 2015;350:g7771.

Berntsen S, Soderstrom-Anttila V, Wennerholm UB, Laivuori H, Loft A, Oldereid NB, Romundstad LB, Bergh C, Pinborg A. The health of children conceived by ART: 'the chicken or the egg?'. *Hum Reprod Update* 2019;25:137–158.

Chen XK, Wen SW, Bottomley J, Smith GN, Leader A, Walker MC. In vitro fertilization is associated with an increased risk for pre-eclampsia. *Hypertens Pregnancy* 2009;28:1–12.

Danhof NA, van Wely M, Repping S, Koks C, Verhoeve HR, de Bruin JP, Verberg MFG, van Hooff MHA, Cohlen BJ, van Heteren CF et al.; SUPER Study Group. Follicle stimulating hormone versus clomiphene citrate in intrauterine insemination for unexplained subfertility: a randomized controlled trial. *Hum Reprod* 2018;33:1866–1874.

De Geyter C, Calhaz-Jorge C, Kupka MS, Wyns C, Mocanu E, Motrenko T, Scaravelli G, Smeenk J, Vidakovic S, Goossens V et al. ART in Europe, 2014: results generated from European registries by ESHRE: the European IVF-monitoring Consortium (EIM) for the European Society of Human Reproduction and Embryology (ESHRE). *Hum Reprod* 2018;33:1586–1601.

Dunietz GL, Holzman C, McKane P, Li C, Boulet SL, Todem D, Kissin DM, Copeland G, Bernson D, Sappenfield WM et al. Assisted reproductive technology and the risk of preterm birth among primiparas. *Fertil Steril* 2015;103:974–979.e1.

Gonzalez-Comadran M, Urresta Avila J, Saavedra Tascon A, Jimenez R, Sola I, Brassesco M, Carreras R, Checa MA. The impact of donor insemination on the risk of preeclampsia: a systematic review

and meta-analysis. *Eur J Obstet Gynecol Reprod Biol* 2014; **182**: 160–166.

Hofteizer L, Hof MHP, Dijks-Elsinga J, Hogeveen M, Hukkelhoven C, van Ling RA. From population reference to national standard: new and improved birthweight charts. *Am J Obstet Gynecol* 2019; **220**:383.e1–383.e17.

Hunault CC, Habbema JD, Eijkemans MJ, Collins JA, Evers JL, Te Velde ER. Two new prediction rules for spontaneous pregnancy leading to live birth among subfertile couples, based on the synthesis of three previous models. *Hum Reprod* 2004; **19**: 2019–2026.

Isaksson R, Gissler M, Tiitinen A. Obstetric outcome among women with unexplained infertility after IVF: a matched case-control study. *Hum Reprod* 2002; **17**:1755–1761.

Joo BS, Park SH, An BM, Kim KS, Moon SE, Moon HS. Serum estradiol levels during controlled ovarian hyperstimulation influence the pregnancy outcome of in vitro fertilization in a concentration-dependent manner. *Fertil Steril* 2010; **93**:442–446.

Kallen B, Finnstrom O, Lindam A, Nilsson E, Nygren K-G, Otterblad Olausson P. Trends in delivery and neonatal outcome after in vitro fertilization in Sweden: data for 25 years. *Hum Reprod* 2010; **25**: 1026–1034.

Lambert RD. Safety issues in assisted reproductive technology: aetiology of health problems in singleton ART babies. *Hum Reprod* 2003; **18**:1987–1991.

Magnus MC, Ghaderi S, Morken N-H, Magnus P, Bente Romundstad L, Skjærven R, Wilcox AJ, Eldevik Håberg S. Vanishing twin syndrome among ART singletons and pregnancy outcomes. *Hum Reprod* 2017; **32**:2298–2304.

Messerlian C, Maclagan L, Basso O. Infertility and the risk of adverse pregnancy outcomes: a systematic review and meta-analysis. *Hum Reprod* 2013; **28**:125–137.

Ombelet W, Martens G, Bruckers L. Pregnant after assisted reproduction: a risk pregnancy is born! 18-years perinatal outcome results from a population-based registry in Flanders. *Facts Views Vis Obgyn* 2016; **8**:193–204.

Pandey S, Shetty A, Hamilton M, Bhattacharya S, Maheshwari A. Obstetric and perinatal outcomes in singleton pregnancies resulting from IVF/ICSI: a systematic review and meta-analysis. *Hum Reprod Update* 2012; **18**:485–503.

Pelinck MJ, Hadders-Algra M, Haadsma ML, Nijhuis WL, Kiewiet SM, Hoek A, Heineman MJ, Middelburg KJ. Is the birthweight of singletons born after IVF reduced by ovarian stimulation or by IVF laboratory procedures? *Reprod Biomed Online* 2010; **21**: 245–251.

Pereira N, Reichman DE, Goldschlag DE, Lekovich JP, Rosenwaks Z. Impact of elevated peak serum estradiol levels during controlled ovarian hyperstimulation on the birth weight of term singletons from fresh IVF-ET cycles. *J Assist Reprod Genet* 2015; **32**:527–532.

Pinborg A, Loft A, Nyboe Andersen A. Neonatal outcome in a Danish national cohort of 8602 children born after in vitro fertilization or intracytoplasmic sperm injection: the role of twin pregnancy. *Acta Obstet Gynecol Scand* 2004; **83**:1071–1078.

Pinborg A, Wennerholm UB, Romundstad LB, Loft A, Aittomaki K, Soderstrom-Anttila V, Nygren KG, Hazekamp J, Bergh C. Why do singletons conceived after assisted reproduction technology have adverse perinatal outcome? Systematic review and meta-analysis. *Hum Reprod Update* 2013; **19**:87–104.

Tjon-Kon-Fat RI, Bensdorp AJ, Scholten I, Repping S, van Wely M, Mol BW, van der Veen F. IUI and IVF for unexplained subfertility: where did we go wrong? *Hum Reprod* 2016; **31**:2665–2667.

van Eekelen R, Tjon-Kon-Fat RI, Bossuyt PMM, van Geloven N, Eijkemans MJC, Bensdorp AJ, van der Veen F, Mol BW, van Wely M. Natural conception rates in couples with unexplained or mild male subfertility scheduled for fertility treatment: a secondary analysis of a randomized controlled trial. *Hum Reprod* 2018; **33**: 919–923.

Wang R, Danhof NA, Tjon-Kon-Fat RI, Eijkemans MJ, Bossuyt PM, Mochtar MH, van der Veen F, Bhattacharya S, Mol BWJ, van Wely M. Interventions for unexplained infertility: a systematic review and network meta-analysis. *Cochrane Database Syst Rev* 2019; **9**: CD012692.