

# Corrigendum. Non-invasive pre-implantation genetic testing of human embryos: an emerging concept

C. Farra<sup>1</sup>, F. Choucair<sup>2</sup>, and J. Awwad<sup>2,\*</sup>

<sup>1</sup>Department of Pathology and Laboratory Medicine, Medical Genetics Unit, American University of Beirut Medical Center, PO Box 11-236, Riad El-Solh, Beirut 1107 2020, Lebanon <sup>2</sup>Department of Obstetrics and Gynecology, Division of Reproductive Endocrinology and Infertility, American University of Beirut Medical Center, PO Box 11-236, Riad El-Solh, Beirut 1107 2020, Lebanon

\*Correspondence address. AUBMC Fertility Center, American University of Beirut Medical Center, PO Box 11-236, Riad El-Solh, Beirut 1107 2020, Lebanon. Tel: +961-1350000 ext. 5606; Fax: +961-1370829; E-mail: jawwad@aub.edu.lb

Hum Reprod 2018;33:2162–2167

The authors of the above article would like to apologise for an error in the 'Clinical studies' section of 'BF sampling'. The article originally reported that the median volume of blastocoel fluid (BF) collected by Palini *et al.* (2013) was 4 µl. The actual volume collected by Palini *et al.* was 0.3–0.5 nl expelled into 4 µl of 1 mmol/l Tris-HCl and 0.1 mmol/l EDTA (pH 8.0).

The electronic version of this article has been updated at <https://doi.org/10.1093/humrep/dey314>. The authors would

like to assure readers that this does not affect any content of the article.

## Reference

Palini S, Galluzzi L, De Stefani S, Bianchi M, Wells D, Magnani M, Bulletti C. Genomic DNA in human blastocoele fluid. *Reprod BioMed Online* 2013; **26**:603–610.